

INDEX TO VOLUME X*

New names, and the final members of new combinations, are in **bold face type**.

- Abies, 198; balsamea, 48; grandis, 9, 99, 100; lasiocarpa, 4, 99
 Acacia Richii, 88
 Acalypha virginica, 169
 Acer neomexicanum, 241
 Achillea lanulosa, 39
 Aecidium Compositarium, 41; micro-punctum, 37
 Agaricaceae of tropical North America—VII, The, 15; VIII, 62
 Agaricus, 16, 25, 29, 30, 67, 68, 69, 70, 81, 85; **angustifolius**, 73, 74; antillarum, 32; arvensis, 77; bambusigenus, 73, 74; bulbillosus, 22; bullaceus, 17; campanulatus, 31; campester, 73, 76, 77, 78; **cinchonensis**, 73, 76; commiscibilis, 19; copriniceps, 21; dichrous, 30; disseminatus, 21, 26; **Earlei**, 74, 79; epibates, 19; euthugrammus, 21; fimicola, 32; fimetarius, 83; fortunatus, 33; guadelupensis, 74, 81; **herradurensis**, 74, 78; hiaseus, 28; hololepis, 69; **Hornei**, 74, 80; junus, 74, 78; **Johnstonii**, 73, 75; minutulus, 26; modestus, 29; **ochraceidiscus**, 74, 80; palmigena, 29; papilionaceus, 31, 32; phalarum, 32; plumiger, 23; plutonius, 30; **praemagnus**, 74, 78; pratensis, 73, 77, 78; pseudotener, 25; Sallei, 81; scatigenus, 17; **Shaferi**, 74, 81; silvicola, 75; solidipes, 31; **subpratensis**, 74, 75; **subsilvicola**, 73, 75; subviridis, 30; tricholepis, 69; **Venus**, 73, 76; **xuchilensis**, 74, 79; yucatanensis, 81
 Agoseris, 242
 Agrimonia striata, 41, 244, 254
 Agropyron Bakeri, 258, 260; Smithii, 251, 262; tenerum, 37, 39, 251
 Allantopora, 240
 Allionia linearis, 247, 248
 Allodus commutata, 35; Douglassii, 35; vertisepta, 35
 Alnus tenuifolia, 239, 241
 Altitudinal range of forest fungi. Notes on, 4
 Amanita, 101; Frostiana, 48; muscaria, 48
 Amanitopsis, 101; volvata, 48
 Amarella heterosepala, 40; strictiflora, 40
 Amorphia canescens, 260
 Anastraphia, 167; Northrupiana, 166
 Andropogon scoparius, 37, 38
 Androsace diffusa, 242
 Anellaria, 30
 Anogra coronopifolia, 258, 260
 Apiospora, 243; sepincoliformis, 243
 Apiosporae, 243
 Apiosporella, 243; **cornina**, 243
 Arabis kochii, 198
 Arbutus Menziesii, 100
 Arenaria Fendleri, 39, 242, 247
 Argentina anserina, 262; argentea, 262
 Armillaria, 266; albolanates, 101; mellea, 10
 Artemisia alba, 38; dracunculoides, 37; franerioides, 40; frigida, 240, 251, 261; gnaphalodes, 40; redolens, 37; scouleriana, 250
 Arthrobotryum (?) pestalozzioides, 263
 Arthur, J. C., Uredinales of Costa Rica based on collections by E. W. D. Holway, 111
 Asclepias speciosa, 254
 Ascochyta **Boutelouae**, 257; graminicola, 257; Miyakei, 286; mori, 285, 286; moricola, 286
 Ascochyta, 258; **agropyrina**, 258
 Ascomycetes, 239
 Ascomycetes and lower fungi from New Mexico, New or noteworthy, 239
 Aster Novae-Angliae, 254; vallicola, 256; Wootonii, 37
 Asterophora Clavus, 97

* It has been considered unnecessary to include here the species listed in the three following articles, since they are already indexed or specially listed.

Arthur: Uredinales of Costa Rica. See p. 150.

Brenckle: North Dakota fungi. See p. 199.

Weir: Altitudinal range of forest fungi. See p. 11.

- Astragalus oreophilus*, 254
Atriplex canescens, 240, 261
Atylospora, 15, 18, 21, 29; **albipes**, 18, 22, 33; **bulbillosa**, 18, 22; **byssina**, 18, 20, 33; **cinchonensis**, 19, 24, 33; **commiscibilis**, 18, 19; **coprinoceps**, 18, 21; **cubensis**, 18, 23, 33; **diminutiva**, 18, 19, 33; **epibates**, 18, 20; **euthugramma**, 18, 21; **fuliginosa**, 18, 25, 33; **lateritia**, 18, 20, 33; **mammillata**, 19, 23, 33; **mexicana**, 18, 21, 33; **Musae**, 18, 21; **pallidispora**, 18, 22, 33; **plana**, 19, 24, 33; **plumigera**, 18, 23; **pseudotenera**, 19, 25; **Roystoneae**, 19, 24; **tigrina**, 18, 19
Azalea mollis, 164
Baccharis Wrightii, 248
Bacterium Juglandis, 99
Betula fontinalis, 251; *occidentalis*, 100
Bjerkandera adusta, 109
Blepharoneuron tricholepis, 251
Boletus luteus, 45
Botrytis, 86; *canescens*, 86; *liliorum*, 86
Bouteloua gracilis, 257
 Brenckle, J. F., North Dakota fungi—II, 199
Bromus ciliatus, 37; *Porteri*, 37
 Bruner, Stephen C., Johnston, John R., and, A *Phyllachora* of the royal palm, 43
 Burlingham, Gertrude S., New species of *Russula* from Massachusetts, 93
Calamagrostis, 167; *hyperborea americana*, 252
Camarosporium Amorphae, 260; *compositarum*, 261; **Estrelti**, 261; *patagonicum*, 261; **wistarianum**, 165; **yuccaesedum**, 261
Campanularius, 16, 29, 30; **anomalus**, 31, 32; *campanulatus*, 31, 32; **solidipes**, 31, 32, 33
Carex, 167; *Douglasii*, 40; *nebraskensis*, 38
Cassiope mertensiana, 4
Castilleja integra, 36, 263; *linariaefolia*, 36; *sulphurac*, 36, 37
Cathartolinum australe, 36
Cenangium, 46; *urceolatum*, 46
Cephaleuros virescens, 291
Cercocarpus montanus, 240, 250, 252
Cercospora montana, 263; *pini-densiflorae*, 89
Ceriospora Dubyi, 244; *montaniensis*, 244, 246
Cerrina unicolor, 110
Chalymota, 30
Chamaecyparis, 189
Chamaesyce serpyllifolia, 41
Cheirinia, 262
Chenopodium album, 259
Chitonia, 85
Chrysomyxa Weirii, 98
Chrysopsis hispida, 247
Chrysothamnus graveolens, 247
Cirsium coloradense, 37; *ochrocentrum*, 37; *polyanthus*, 41; *Richardsoni*, 41
Citrus, 88
Cladosporium fasciculatum, 263; *herbarum*, 262; *herbarum cerealium*, 263
Clarkeinda, 16
Clathrospora permunda, 248
Clavaria, 57; *ardenia*, 54, 55, 57; *brachiata*, 56; *contorta*, 53, 55, 56, 57; *fistulosa*, 53, 54, 55, 56, 57; *inaequalis*, 53; *junceae*, 53, 54, 56, 57; *junceae vivipara*, 56; *macrorrhiza*, 54, 55, 57; *vermicularis*, 53
Clavaria fistulosa group, The, 53
Clematis, 166; *calycina*, 166; *integri-folia*, 166; *ligusticifolia*, 38, 244, 251, 264; *paniculata*, 165, 166; *vitalba*, 166
Clitocybe farinacea, 180; *illudens*, 292
Coccomyces Kerriae, 99
Coleosanthus grandiflorus, 39; *reniformis*, 246
Coleosporium ribicola, 35
 Collections by E. W. D. Holway
Uredinales of Costa Rica based on, 111
Collema, 236, 238; *pulposum*, 235, 236, 238
Collemaceae, A new genus and species of the, 235
Collemodes, 236; **Bachmanianum**, 237, 238
Collybia conigena, 55; *esculenta*, 55; *velutipes*, 266
Comandra pallida, 254
Compositae, 247
 Conidial formation in *Sphaeronomia fimbriatum*, 155
Coniothyrium concentricum Yuccaglaucac, 257; *myriocarpum*, 257; *olivaceum Salsolae*, 257; *olivaceum Thermopsisidis*, 257; *sepium*, 256
Convolvulus sepium, 256
Coprinopsis, 82
Coprinus, 19, 21, 28, 82, 84; *armillaris*, 82, 83; **cinchonensis**, 82, 85; *cubensis*, 82, 83; *fimetarius*, 82, 83; *jalapensis*, 82, 83; *jamaicensis*, 82, 84; *mexicanus*, 82, 84; *mica-*

- ceus, 28, 83; plicatilis, 82, 85;
Spraguei, 82, 85
Coriollus serialis, 290
Corioliopsis rigida, 269
Coriolus prolificans, 110; pubescens,
290
Cornu instolonea, 254, 255
Corticum, 239
Cortinarius, 15, 46; corrugatus, 234;
salmonicolor, 46
Cortinopsis, 62
Costa Rica based on collections by E.
W. D. Holway, Uredinales of, 111
Crataegus erythropoda, 36, 252, 253
Crepidotus, 16; alveolatus, 16;
fumosifolius, 16; musicola, 16
Cronartium coleosporoides, 36, 37
Cryptostictis **utensis**, 260
Cucurbitaria Ribis, 250; Rosae, 250
Cultures with Melampsorae on Popu-
lus, 194
Cup-fungi. Photographs and descrip-
tions of.—VII. The genus Under-
woodia, 1
Cycloporus Greenei, 47
Cylindrocladium scoparium, 99
Cyrtorhynca ranunculina, 253
Cytodiplopora mori, 287
Cytospora chrysosperma, 266

Daedalea confragosa, 99; unicolor,
99, 110
Daldinia, 279, 280, 283, 284; aspera,
280; cingulata, 280; concentrica,
280, 281, 282, 283, 284; durissima,
280; loculata, 280; vernicosa, 277,
278, 279, 280, 281, 282, 283
Daldinia vernicosa—a pyroxylophi-
lous fungus, 277
Danthonia, 252; intermedia, 248, 252
Dasiophora fruticosa, 36
Dasyphypha Willkommii, 9
Dasystephana Bigelovii, 38
Deconica, 15, 17; bullacea, 17, 18;
scatigena, 17
Delphinium robustum, 245
Dematiaceae, 262
Dematium pullulans, 263
Descriptions of cup-fungi. Photo-
graphs and.—VII. The genus Under-
woodia, 1
Deuteromycetes, 253
Diaporthe oligocarpoides, 244
Diatrype albobrunnea, 241; cornuta,
241; **Standleyi**, 240
Diatrypella Placenta, 241
Dictyochora **Gambellii**, 166
Didymella **Eurotiae**, 243; **nigrescens**,
242; **nigrificans**, 243
Didymosphaeria, 244
Diervilla Diervilla, 164, 165
Dimerosporium mori, 288
Diplodia, 45; hyalospora, 165
Distribution of fungi in Porto Rico,
The, 58
Dodge, B. O., Studies in the genus
Gymnosporangium—III. The ori-
gin of the teleutospore, 182
Dothichiza populea, 170
Dothidea Haydeni, 256
Dothideaceae, 251
Dothideales, 243
Dothidella insculpta, 251
Dothiorella, 255; **phomopsis**, 255
Drosophila, 16, 18, 21, 62, 67; ap-
pendiculata, 62, 63, 64, 66, 71; **atri-
castanea**, 63, 66; **brevipes**, 62, 63;
caespitosa, 63, 67; **campestris**, 63,
64; **castaneidisca**, 62, 63; **flocu-
losa**, 63, 64; **jalapensis**, 63, 65;
pallidospora, 63, 64; **tenuis**, 63, 65;
tepeitensis, 63, 65; **truncatispora**,
63, 66
Dugaldea Hoopesii, 41

Echinochloa zelayensis, 41
Echinodontium tinctorium, 5, 7, 10
Elymus canadensis, 38, 245
Entoloma, 30
Epilobium adenocladon, 263; novo-
mexicanum, 41
Eriogonum alatum, 259; Jamesii, 40;
racemosum, 40
Erysibaceae, 239
Eurotia lanata, 243, 254
Euryachora betulina, 251
Eutypella Brunaudiana **Ribis-aurei**,
240; herbicola, 240

Fabraea litigiosa, 253
Fairman, Charles E., New or note-
worthy Ascomycetes and lower
fungi from New Mexico, 239;
Notes on new species of fungi
from various localities—II, 164
Fink, Bruce, A new genus and species
of the Collemaceae, 235; The dis-
tribution of fungi in Porto Rico,
58
Firmiana platanifolia, 90
Flocculosa, 233
Fomes annosus, 10; lucidus, 45, 290;
officinalis, 269; pini, 5, 7, 10; un-
gulatus, 100
Forest fungi, Notes on the altitudinal
range of, 4
Fungi from New Mexico, New or
noteworthy Ascomycetes and lower,
239
Fungi from various localities—II,
Notes on new species of, 164
Fungi, Illustrations of—XXVIII,
107; XXIX, 177

- Fungi in Porto Rico, The distribution of, 58
 Fungi—II, North Dakota, 199
 Fungi, Notes and translations—, New Japanese, IV, 86; V, 285
 Fungi, Notes on the altitudinal range of forest, 4
 Fungus, *Daldinia vernicosa*—a pyroxylophilous, 277
Fusarium, 267
Fusicladium Cerasi, 263
Fusicoccum putrefaciens, 46

Gaertneria acanthicarpa, 254
Galium boreale, 241, 258
Ganoderma sessile, 290
Gaultheria humifusa, 4
Gaura induta, 261, 262; *parviflora*, 261
Gautieria, 266; *plumbea*, 266
 Genus and species of the Collema-ceae, A new, 235
 Genus *Gymnosporangium*—III. The origin of the teleutospore, Studies in the, 182
Geophila, 70
Gibberidia arthrophytha, 246
Globifomes graveolens, 267
Glochidion obovatum, 88
Gleophyllum trabeum, 290
Gloeosporium Potentillae, 262; *venetum*, 46
Glyptosperma, 62
Gomphidius, 16, 69; *jamaicensis*, 69
Grindelia aphanactis, 41
Grossularia inermis, 35; *leptantha*, 241, 250
Gymnochilus, 62; *caespitosus*, 67; *campestris*, 64; *flocculosus*, 64; *Musae*, 21; *Roystoniae*, 24
Gymnolomia multiflora, 37
Gymnopus, 177
Gymnosporangium, 182, 189, 191; *Betheli*, 36; *clavariaeforme*, 182, 183, 189, 190, 192; *clavipes*, 183, 190; *fraternum*, 183; *globosum*, 183, 187, 189, 192; *juniperi-virginianae*, 183; *macropus*, 182, 183, 184, 187, 191; *nidus-avis*, 190, 192; *Sabinae*, 182; *transformans*, 183
Gymnosporangium—III. The origin of the teleutospore, Studies in the genus, 182

 Harper, Edward T., *Hypoloma aggregatum* and *H. delineatum*, 231; The *Clavaria fistulosa* group, 53
Hedysarum pabulare, 40
Helianthus annuus, 41, 261; *Maximiliani*, 246, 251
Helicobasidium tanakae, 89, 90, 91
Heliopsis scabra, 245, 255

Helvella crispa, 2
Hendersonia Agropyri, 260; *calycina*, 166; *Clematidis*, 166; *crastophila*, 260; *Eriogoni*, 259; *foliorum*, 260; *hortilecta*, 165; *Leucelene*, 249; *Petalostemonis*, 249, 260; *Rubi Clematidis*, 166; *sarmentorum Clematidis*, 166; *Stanleyellae*, 259; *subcultriformis*, 260
Heracleum lanatum, 254
Herpotrichia nigra, 6
Heuchera parvifolia, 39, 240
Hicoria alba, 279
Hieracium Fendleri, 38
 Holway, Uredinales of Costa Rica based on collections by E. W. D., 111
Hordeum trifurcatum, 42
Houstonia minor, 168
 Hubert, Ernest E., Weir, James R., and, Cultures with *Melampsora* on *Populus*, 194
Humulus americanus, 244, 259
Hura crepitans, 68
Hyalodidymae, 242, 257
Hyalophragmiae, 259
Hyalosporae, 241, 253
Hydnum, 110
Hymenochaete noxia, 45
Hypholoma, 16, 67, 68, 71, 84, 234; *aggregatum*, 231, 232, 233; *aggregatum sericeum*, 232, 233; *bermudiense*, 72; *caespitosum*, 67; *campestre*, 64; *delineatum*, 231, 232, 233, 234; *echiniceps*, 232, 233; *fasciculare*, 67, 68; *flavovirens*, 67, 68; *flocculosum*, 64; *hypoxanthum*, 232; *lacrimabundum*, 231, 232, 233, 234; *papillatum*, 67; *populinum*, 232, 233; *Pseudostorea*, 232, 233; *rugoccephalum*, 234; *silvestre*, 233; *Storea*, 231, 232, 233; *Storea caespitosum*, 231, 232, 233; *tuberculatum*, 67, 68; *velutinum*, 232, 234
Hypholoma aggregatum and *H. delineatum*, 231
Hypholoma delineatum, *Hypholoma aggregatum* and, 231
Hypholomopsis, 62
Hypocreaceae, 251
Hypoxylon cerebrinum, 280, 284
Hysteriaceae, 252
Hysterium Notarisianum, 252; *Standleyanum*, 252
Hysterographium Bakeri, 252

 Illustrations of fungi—XXVIII, 107; XXIX, 177
 Index to American mycological literature, 49, 102, 172, 226, 271, 295
Incocybe, 15, 222
Ionoxalis violacea, 41

- Ipomoea batatas*, 155
Iris missouriensis, 242
Irpey, 110
Isocoma heterophylla, 247, 254, 261
 Japanese fungi, Notes and translations—, New, IV, 86; V, 285
 Johnston, John R., and Bruner, Stephen C., *A Phyllachora* of the royal palm, 43
Juglans, 90
Juncus balticus, 263
Juniperus scopulorum, 36
Kalmia latifolia, 268
Kellermannia Sisyrrinchii, 259; *yucca*-*caegena*, 259
Kerria japonica, 90, 99
Koeleria cristata, 252, 263
Kuhnia rosmarinifolia, 38, 247
Laccaria amethystea, 178; *laccata*, 178, 179; *striatula*, 179
Lachnella flammea, 252; *rhoina*, 252
Lachrymaria, 62
Laciniaria punctata, 247
Lactaria Allardii, 265; *coleopteris*, 265; *Curtisii*, 265; *furcatus*, 265; *lentus*, 265; *subplinthogalus*, 265; *subterminosus*, 265
Lactuca pulchella, 41, 262
Larix, 194, 195, 196, 198; *europaea*, 197; *laricina*, 195; *lyallii*, 5, 197, 198; *occidentalis*, 194, 195, 196, 197
Lathyrus arizonicus, 40; *decaphyllus*, 40; *leucanthus*, 40
Lavauxia flava, 262
Ledum glandulosum, 4
 Lehman, S. G., *Conidial formation in Sphaeronema fimbriatum*, 155
Lentinus lepideus, 8
Lentodium tigrinum, 290
Lenzites sepiaria, 8
Lepiota, 75, 81, 82, 83; *cretacea*, 82
Leptonia conica, 178
Leptoniella conica, 178
Leptosphaeria Coleosanthi, 246; *culmifraga minuscula*, 245; *Doliolum*, 245; *dumetorum*, 244; *Helianthi*, 246; *lupincola*, 245; *nigricans*, 245; *nigricans Grindeliae*, 245; *ogilviensis*, 245; *praeclara typhiseda*, 245; *Quamoclidii*, 246; *rubrotincta*, 246; *Senecionis*, 248; *tenera*, 245
Leucelene arenosa, 249, 250
Ligusticum Porteri, 246
Lilium longiflorum, 86
Linum Lewisii, 36
Lithospermum multiflorum, 247
Locellina, 15
Lophidiopsis nuculoides, 252
Lophiosphaera perpusilla, 167
Lophiostoma collinum, 167; *quadri-nucleatum*, 251
Lophiostomataceae, 251
Lophiotrema Mollerianum, 167; *pusillum*, 167; *stenogramma*, 167
Lophodermium arundinaceum, 252
 Lower fungi from New Mexico, New or noteworthy *Ascomycetes* and, 239
Lupinus ingratus, 245
Lycoperdon giganteum, 292
Lycurus phleoides, 248
Lygodesmia juncea, 262
Machaeranthera Bigeloviae, 245; *Bigelowii*, 263
Macrophoma cornina, 255; *minuta*, 285
Macrosporium commune, 263
Malvastrum coccineum, 39
Marasmius, 177; *dichrous*, 180; *insititius*, 181
Massachusetts, New species of *Russula* from, 93
Massaria, 240
Medicago sativa, 253, 254
Melampsora albertensis, 41, 194, 195, 196, 197, 198; *Bigelovii*, 36; *Lini*, 36; *medusae*, 194, 195, 196, 197, 198; *occidentalis*, 98
Melampsorae, 194, 198
Melampsorae on *Populus*, Cultures with, 194
Melamporella elatina, 198
Melanconiaceae, 262
Melanconiales, 262
Melanopsamma pomiformis, 244
Melanotus, 15, 16; *fumosifolius*, 16; *musicola*, 16
Melia Azedarach, 88
Melilotus alba, 244
Meliola, 288
Mertensia caelestina, 242, 248
Metasphaeria Senecionis, 248
Micranthes arguta, 39
Microdiplodia Anograe, 258; *Dier-villae*, 165; *galiicola*, 241, 258; *Leucelenes*, 249; *Viciae*, 258
Micromphale ulmarium, 290
Monarda comata, 38; *stricta*, 38
Moniliales, 262
Morus, 90; *alba*, 87, 91, 92, 285, 286, 287, 288; *alba Nezumigaeshi*, 288
Muhlenbergia cuspidata, 251; *trifida*, 39, 251
 Murrill, W. A., Illustrations of fungi—XXVIII, 107; XXIX, 177; The *Agaricaceae* of Tropical North America—VII, 15; VIII, 62
Mycena galericulata, 179; *viscidipes*, 177

- Mycosphaerella, 258; Iridis, 242;
 Oenotherae, 258; pachyasca, 242;
 Primulae, 242; tingens, 242
 Myriangaceae, 46
- Naematoloma, 67
 Naucoria, 21; coprinoceps, 21;
 euthugramma, 21
 Neopeckia coulteri, 6
 New and noteworthy species, Studies in North American Peronosporales—VII, 168
 New genus and species of the Collemaeae, A, 235
 New Japanese fungi, Notes and translations—IV, 86; V, 285
 New Mexico in 1916, Rusts and smuts collected in, 34
 New Mexico, New or noteworthy Ascomycetes and lower fungi from, 239
 New or noteworthy Ascomycetes and lower fungi from New Mexico, 239
 New species of fungi from various localities—II, Notes on, 164
 New species of Russula from Massachusetts, 93
 North America—, The Agaricaceae of tropical, VII, 15; VIII, 62
 North American Peronosporales—VII, New and noteworthy species, Studies in, 168
 North Dakota fungi—II, 199
 Nostoc, 235
 Notes and brief articles, 45, 97, 170, 222, 265, 289
 Notes and translations—, New Japanese fungi, IV, 86; V, 285
 Notes on new species of fungi from various localities—II, 164
 Notes on the altitudinal range of forest fungi, 4
 Noteworthy Ascomycetes and lower fungi from New Mexico, New or, 239
 Noteworthy species, Studies in North American Peronosporales—VII, New and, 168
 Nothopatella moricola, 91
 Nummularia Bulliardii, 279
 Nuttalia Rusbyi, 244
- Odostemon repens, 41
 Oenothera Hookeri, 262
 Omphalia fibula, 179
 Omphalopsis fibula, 179
 Ophiobolus claviger, 250; collapsus, 250
 Ophiodothia Haydeni, 256
 Oreobatus deliciosus, 37
 Origin of the teleospore, Studies in the genus Gymnosporangium—III, The, 182
 Osmunda cinnamomea, 94; regalis, 94
 Otthia Clematidis, 244; fruticola, 244
 Oxytropis deflexa, 41; Lambertii, 262
- Padus melanocarpa, 263
 Palm, A Phyllachora of the royal, 43
 Paneolus, 30; campanulatus, 31; fimicola, 32; papilionaceus, 32; phalenarum, 32; solidipes, 31, 33
 Panicum virgatum, 40
 Patella **oreophila**, 253
 Paulownia, 92; tomentosa, 90
 Pedicularis fluviatilis, 38, 254; Grayi, 36
 Pentstemon Torreyi, 37
 Pericome caudata, 245
 Peridermium coloradense, 198; ribicola, 35
 Peronospora, 168, 169; grisea, 168; Polygoni, 168; Seymourii, 168
 Peronosporales—VII, New and noteworthy species, Studies in North American, 168
 Pestalozzia, 264
 Petalostemon oligophyllus, 249, 260
 Pezizales, 252
 Phaeoapiospora, 243
 Phaeoapiosporae, 244
 Phaeodictyae, 247, 260
 Phaeodidymae, 244, 258
 Phaeolus sistotremonoides, 290
 Phaeophragmiae, 244, 259
 Phaeosporae, 241, 256
 Philocopa coeruleotecta, 222
 Phloeospora Oxytropidis, 262
 Phlox, 35
 Pholiota, 15, 71, 98; adiposa, 98; erebia, 234; flammans, 98; squarrosa, 98
 Phoma, 241, 243, 245, 247, 250, 255, 266; asclepiadea, 254; Astragali, 254; complanata, 254; Corni-Sueciae, 254; **Estrelti**, 254; exigua, 254; herbarum, 250, 254; herbarum Medicaginis, 254; herbarum Solidaginis, 255; lupicola, 246; Mororum, 87; oleracea, 255; Rudbeckiae, 255; **Sidalceae**, 255; thalietrina, 255; **verbascicarpa**, 164
 Phomopsis **ericaceana**, 164; Kalmiae, 268
 Photographs and descriptions of cup-fungi—VII, The genus Underwoodia, 1
 Phragmidium Andersoni, 36; imitans, 41; montivagum, 37; Peckianum, 37; Potentillae, 41
 Phyllachora, 43; Ambrosiae, 251;

- Blepharoneuri**, 251; *Haydeni*, 256;
Roystoneae, 43, 44; *Trifolii*, 251;
vulgata, 251
Phyllachora of the royal palm, A, 43
Phyllactinia corylea, 239
Phyllodoce empetrifomis, 4
Phyllosticta Crataegi, 253; *Kuwacola*,
87
Phyllostictaceae, 253
Physalospora Ambrosiae, 251; *Ar-*
thuriana, 251; *Galii*, 241; *minuta*,
285
Picea engelmanni, 5, 198
Pilosace, 16, 68; *hololepis*, 69; *tri-*
cholepis, 69
Pinus albicaulis, 4; *cembra*, 5; *con-*
torta, 5; *densiflora*, 89; *edulis*, 35;
flexilis, 5; *pungens*, 48; *rigida*, 48;
Strobilus, 48
Pittosporum undulatum, 90
Placosphaerella, 256
Placosphaeria decipiens, 256
Plasmopara Acalyphae, 169
Platystomum compressum, 252; **phyl-**
logenium, 166
Plectodiscella veneta, 46
Pleospora, 243; *Bardanae*, 247; *colo-*
radensis, 248; *compositarum*, 247;
herbarum, 246, 247, 248; *herbarum*
microspora, 247; *infectoria*, 248;
rubicunda, 248; *Senecionis*, 248;
vulgaris, 248; *vulgatissima*, 248
Pluteopsis, 18
Poa Bigelovii, 38; *pratensis*, 38
Poinsettia dentata, 41
Polemonium confertum, 248
Polygonum sawatchense, 254
Polyporus, 108; *adustus*, 109; *albel-*
lus, 99; *amorphus*, 48, 109; *bru-*
malis, 108; *elegans*, 100; *glomerat-*
us, 170; *irregularis*, 109; *leuco-*
spongia, 7, 8; *Polyporus*, 108;
Schweinitzii, 10, 269
Polystictus conchifer, 108; *hirsutus*, 8
Polythrincium Trifolii, 263
Populus, 194, 198; *angustifolia*, 241,
244, 250, 257; *aurea*, 41; *deltoides*,
195; *tremuloides*, 194, 195, 196,
197; *trichocarpa*, 194, 195, 196, 197
Populus, Cultures with *Melampsorae*
on, 194
Poronidulus conchifer, 108
Porto Rico, The distribution of fungi
in, 58
Potentilla filipes, 247; *strigosa*, 41
Pratella, 73
Primula angustifolia, 260
Prunulus, 20, 177; *galericulatus*, 179;
viscidipes, 177
Prunus americana, 263; *Armeniaca*
Ansu, 90; *donarium*, 90; *Mume*,
90; *Persica*, 88; *salicina*, 88, 90
Psalliota, 73
Psathura, 18
Psathyra, 18; *albipes*, 33; *bulbillosa*,
22; *hyssina*, 33; *cinchonensis*, 33;
commiscibilis, 19; *cubensis*, 33;
cubispora, 30; *diminutiva*, 33; *epi-*
bates, 20; *fuliginosa*, 33; *lateritia*,
33; *mammillata*, 33; *mexicana*, 33;
pallidispora, 33; *plana*, 33; *plumi-*
gera, 23; *pseudotenera*, 25; *tigrina*,
19
Psathyrella, 15, 18, 25, 28; *arata*, 28;
cubensis, 25, 27; *disseminata*, 26;
Earlei, 25, 27; **grisea**, 25, 26; *hi-*
ascens, 28; **mexicana**, 25, 26;
minutula, 25, 26; *modesta*, 29;
prona, 26; **Stevensonii**, 25, 28
Pseudocymopterus montanus, 39;
multifidus, 39
Pseudopeziza Medicaginis, 253
Pseudotsuga, 194, 195, 196, 198;
macrocarpa, 197, 198; *mucronata*,
196; *taxifolia*, 5, 194, 195, 196, 197
Psilocybe, 15, 17, 18, 29, 33; *antil-*
larum, 32; *antillarum praelonga*,
33; *dichroma*, 29, 30; *fortunata*,
33; *palmigena*, 29; *plutonica*, 29,
30; **orizabensis**, 29; *subviridis*, 30,
68
Psoralea tenuiflora, 248
Pteris aquilina, 94
Puccinia, *Absinthii*, 37; *aemulans*,
37; *Andropogonis*, 36, 37, 38;
Artemisiae, 37; *Asteris*, 37; *Cirsii*,
37; *Clematidis*, 37; *Clintonii*, 38;
conferta, 38; *Ellisiana*, 37, 38; *epi-*
phylla, 38; *Gentianae*, 38; *Grin-*
deliae, 41; *Grossulariae*, 38; *Heli-*
anthi, 41; *hemispherica*, 41; *Hie-*
racii, 38; *Kuhniae*, 38; *Menthae*, 38;
Millefolii, 39; *monoica*, 198; *Mu-*
hlenbergiae, 39; *poculiformis*, 39;
Pseudocymopteri, 39; *Oxalidis*, 41;
Saxifragae, 39; *Sherardiana*, 39;
subdecora, 39; *substerilis*, 39; *Ta-*
raxaci, 41; *tardissima*, 39; *tuber-*
culans, 39; *universalis*, 40; *verti-*
septa, 35; *violae*, 40
Pucciniastrum Agrimoniae, 41; *pust-*
ulatum, 41
Pycnoporus cinnabarinus, 107; *sangui-*
neus, 107
Pyrenophora, 249; *chrysospora po-*
laris, 248; *comata*, 249; **Leucelenes**,
249
Pyronema, 277
Pyropolyporus conchatus, 290
Pyroxylophilous fungus, *Daldinia*
vernica—a, 277
Pyrus Malus, 90; *sinensis*, 90
Quamoclidion multiflorum, 246

- Quercus*, 167; *alba*, 279, 284; *Fendleri*, 241, 250, 252
- Range of forest fungi, Notes on the altitudinal, 4
- Ratibida columnifera*, 244
- Rhabdospora*, 241, 244; **dumetorum**, 245; **gauracea**, 262; **translucens**, 165
- Rhizopogon*, 222
- Rhoads, A. S., *Daldinia vernicosa*—a pyroxyphilous fungus, 277
- Rhododendron albiflorum*, 4
- Rhus*, 250
- Rhysotheca Acalyphae*, 169; *australis*, 169; *illinoisensis*, 169
- Rhytisma salicinum*, 253
- Ribes aureum*, 35, 36, 240, 250; *Grossularia*, 90; *inebrians*, 35, 36, 251, 252; *longiflorum*, 36; *longifolium*, 36; *mesacalerium*, 36; *Wolfii*, 35
- Robillarda Cavarae*, 287; *Celtidis*, 287; *mori*, 287
- Rosa*, 242, 243, 244, 250; *Fendleri*, 37; *Maximiliani*, 37
- Rosellinia*, 45, 98; *parasitica*, 241; *pulveracea*, 241, 257; *Rosarum*, 242
- Royal palm, A *Phyllachora* of the, 43
- Roystonea regia*, 43, 44
- Rubus arizonicus*, 41
- Rudbeckia laciniata*, 41, 255
- Russula*, 93, 95, 267; *alutacea*, 94; *cinerascens*, 267; **Davisii**, 93; **disparalis**, 94; *elatior*, 95; *fallax*, 93; *fragiliformis*, 93, 96; *fragilis*, 93; *glauca*, 93; *heterophylla*, 93; *insignis*, 93; *integra*, 93; *Linnaei*, 96; *magna*, 267; *paludosa*, 95; *pectinatoides*, 93; **perplexa**, 96; **pulchra**, 95, 96; *pungens*, 267; *purpurina*, 96; *rubrotincta*, 95; *subvelutina*, 96; *uncialis*, 96; *vernosa*, 93
- Russula* from Massachusetts, New species of, 93
- Rusts and smuts collected in New Mexico in 1916, 34
- Salix*, 36, 90; *ballotaeflora*, 35; *Bebbiana*, 253; *cordata Watsoni*, 36, 252, 263; *glandulosa Warburgii*, 88; *Scouleriana*, 36; *subcaerulea*, 36; *Wrightii*, 36
- Salsola Pestifer*, 257
- Schizophyllum commune*, 7, 100
- Schizoxylon insigne*, 253
- Schmaltzia Bakeri*, 250, 252
- Sclerotinia parasitica*, 86
- Scolecosporeae*, 250, 261
- Seaver, Fred J., Photographs and descriptions of cup-fungi—VII, The genus *Underwoodia*, 1
- Sebacina spongiosa*, 98
- Senecio amplexens*, 248; *scopolina*, 244; *spartioides*, 254
- Septobasidium*, 89; *acaciae*, 88; *pedicellatum*, 90
- Septoria gaurina*, 261, 262; *Helianthi*, 261; *Oenotherae*, 262; *smilacina*, 262
- Sidalcea neomexicana*, 255
- Sideranthus spinulosus*, 39
- Sisyrinchium demissum*, 259
- Sitanion longifolium*, 42, 262
- Smuts collected in New Mexico in 1916, Rusts and, 34
- Solidago Pitcheri*, 255
- Sparassis radicata*, 100
- Species of the *Collema*ceae, A new genus and, 235
- Species of fungi from various localities—II, Notes on new, 164
- Species, Studies in North American Peronosporales—VII, New and noteworthy, 168
- Spermoedia clavus*, 251
- Sphaeria vernicosa*, 283, 284
- Sphaeriaceae*, 240
- Sphaeriales*, 239
- Sphaeronema*, 155, 161, 162; *fimbriatum*, 155, 156, 160, 162
- Sphaeronema fimbriatum*, Conidial formation in, 155
- Sphaeropsidales*, 253
- Sphaeropsis Diervillae*, 164; **wistariana**, 164
- Sphaerotheca Humuli*, 240
- Sporobolus auriculatus*, 248, 259
- Stagonospora Chenopodii*, 259; *graminella*, 259; **Humuli-americanii**, 259; *mori*, 286
- Standley, Paul C., Rusts and smuts collected in New Mexico in 1916, 34
- Stanleyella Wrightii*, 259, 263
- Stictidiaceae, 45
- Stilbaceae, 263
- Stilbum*, 264
- Stipa Scribneri*, 39; *Vaseyi*, 39, 42
- Strickeria Cercocarpi*, 250; *rhoina*, 250
- Stropharia*, 16, 68, 70, 71; *aeruginosa*, 72; *ambigua*, 101; **bermudiensis**, 70, 72; **caespitosa**, 70, 71; *cubensis*, 70, 72; *floccosa*, 70, 71; *melsperma*, 73; *semiglobata*, 70, 72, 73; *stercoraria*, 73; **troyana**, 70
- Studies in North American Peronosporales—VII, New and noteworthy species, 168
- Studies in the genus *Gymnosporan-*

- gium—III, The origin of the teleutospore, 182
Stypinella Tanakae, 89
Symphoricarpos oreophilus, 242, 252, 253
 Tanaka, Tyôzaburô, New Japanese fungi, Notes and translations—IV, 86; V, 285
Taraxacum taraxacum, 41
Tecoma radicans, 165
Teichospora Cercocarpi, 250; obducens, 250; pygmaea, 250; rhoina, 250; rhypodes, 250; stenocarpa, 250
 Teleutospore, Studies in the genus *Gymnosporangium*—III, The origin of the, 182
Thalictrum dasycarpum, 255; *Fendleri*, 255
Thea sinensis, 88, 90
Thermopsis Pinetorum, 254, 257
Thielavia, 155, 156, 160, 161, 162; *basicola*, 155, 156, 160
Thyridaria tarda, 45
Thyridium, 240; *cingulatum*, 240
Trametes carnea, 100; *cinnabarina*, 107; *hispida*, 100; *serialis*, 100; *variiformis*, 100
 Translations—, New Japanese fungi, Notes and, IV, 86; V, 285
Trifolium Fendleri, 251, 263; *nanum*, 247; *stenolobum*, 247
Trimmatostroma Salicis, 263
Triticum aestivum, 39
 Tropical North America—, The Agaricaceae of, VII, 15; VIII, 62
Tsuga heterophylla, 9, 100, 197; *mertensiana*, 4; *taxifolia*, 100
Tubercularia vulgaris, 263
 Tuberculariaceae, 263
Typha latifolia, 245, 248
Typhula, 57
Tyromyces amorphus, 109
 Uncinula flexuosa, 239
 Underwoodia, 1; *columnaris*, 2
 Uredinales, 35
 Uredinales of Costa Rica Based on collections by E. W. D. Holway, 111
Uromyces Astragali, 41; *Eriogoni*, 40; *Fabae*, 40; *Gentianae*, 40; *graminicola*, 40; *Hedysari-obscuri*, 40; *intricatus*, 40; *proeminens*, 41; *punctatus*, 41; *Rudbeckiae*, 41
Uropyxis sanguinea, 41
 Ustilaginales, 41
Ustilago bromivora, 41; *Crus-galli*, 41; *Hordei*, 42; *hypodytes*, 42; *levis*, 42
Ustilina microspora, 92; *mori*, 91; *zonata*, 45, 46
Vaccinium microphyllum, 4
Vagnera stellata, 262
Valeriana, 35
Valsa Paulowniae, 92
Velutina, 233
Venenarius, 101; *pantherinoides*, 280; *pregammatus*, 101; *umbrinidiscus*, 101
Verbasum Blattaria, 164
Verbena Macdougallii, 250
Veronica, 168; *arvensis*, 168; *perigrina*, 168
Vicia americana, 40, 258
Viola pedatifida, 38, 40
Viorna Scottii, 255
Vitis, 90
 Weir, James R., Notes on the altitudinal range of forest fungi, 4; Hubert, Ernest E., and, Cultures with *Melampsorae* on *Populus*, 194
 Wilson, Guy West, Studies in North American Peronosporales—VII, New and noteworthy species, 168
Xanthoxylum, 90
Xylaria Hypoxylon, 45
Yucca baccata, 261; *glauca*, 257, 259
Zea Mays, 166